



PHYSICAL CHEMISTRY 2012

11th International Conference
on Fundamental and Applied Aspects of
Physical Chemistry

Under the auspices of the
University of Belgrade

Proceedings

The Conference is dedicated to
Professor Ivan Draganić

September 24-28, 2012
Belgrade, Serbia

ISBN 978-86-82475-27-9 Volume I
ISBN 978-86-82475-28-6 Volume II

Title: PHYSICAL CHEMISTRY 2012 (Proceedings)

Editors: S. Anić and Ž. Čupić

Published by: Society of Physical Chemists of Serbia, Studenski trg 12-16,
11158, Belgrade, Serbia

Publisher: Society of Physical Chemists of Serbia

For Publisher: S. Anić, President of Society of Physical Chemists of Serbia

Printed by: “Jovan” Priting and Publishing Company; 200 Copies;

Number of pages: 6+ 497; **Format:** B5; Printing finished in September
2012.

Text and Layout: “Jovan”

200- *Copy printing*

ORGANIZERS

PHYSICAL CHEMISTRY 2012

*11th International Conference on
Fundamental and Applied Aspects of
Physical Chemistry*

*Under the auspices of the
University of Belgrade*

*Organized by
The Society of Physical Chemists of
Serbia*

*in co-operation
with*

Institute of Catalysis Bulgarian Academy of Sciences

*Boreskov Institute of Catalysis of Siberian Branch of the Russian Academy
of Sciences*

Faculty of Physical Chemistry, University of Belgrade, Serbia

*Institute of Chemistry Technology and Metallurgy,
University of Belgrade, Serbia*

Vinča Institute, University of Belgrade, Serbia

Institute of General and Physical Chemistry, Serbia

Faculty of Pharmacy, University of Belgrade, Serbia

COMMITTEES

International Organizing Committee

Chairman: S. Anić (Serbia)

Vice-chairmans: M. Gabrovska (Bulgaria)

V. A. Sadykov (Russia)

B. Adnadjević (Serbia)

Members: Z. Cherkezova-Zheleva (Bulgaria), N. Cvjetićanin (Serbia), S. N. Blagojević (Serbia), M. Daković (Serbia), T. Grozdić (Serbia), Lj. Ignjatović (Serbia), D. Jovanović (Serbia), N. Jović-Jovičić (Serbia), M. Kuzmanović (Serbia), D. Marković (Serbia), N. Kovačević (Serbia), J. Marković-Dimitrić (Serbia), B. Milosavljević (USA), Š. Miljanić (Serbia), N. Miljević (Serbia), A. Milutinović-Nikolić (Serbia), M. Mojović (Serbia), Z. Mojović (Serbia), N. Ostrovski (Serbia), I. Pašti (Serbia), C. Pona (Italy), B. Simonović (Serbia), D. Stanisavljev (Serbia), B. Šljukić (Serbia), N. Vukelić (Serbia), V. Vukojević (Sweden)

International Scientific Committee

Chairman: Ž. Čupić (Serbia)

Vice-chairmans: V. N. Parmon (Russia)

S. Rakovsky (Bulgaria)

Members: A. Antić-Jovanović (Serbia), G. Bačić (Serbia), R. Cervellati (Italy), R. Compton (UK), V. Dondur (Serbia), Biljana Gaković (Serbia), V. Gaspar (Hungary), S. Gorodsky (Russia), V. Goussev (France), K. Hedrih (Serbia), Florou Heleny (Greece), M. Jeremić (Serbia), A. L. Kawczyński (Poland), Lj. Kolar-Anić (Serbia), V. Kuntić (Serbia), S. Mentus (Serbia), S. Milonjić (Serbia), Š. Miljanić (Serbia), Lj. Morozova-Roche (Sweden), P. Misaelides (Greece) M. Perić (Serbia), M. Plavšić (Serbia), V. Sadykov (Russia), G. Schmitz (Belgium), I. Schreiber (Czech), P. Sevcik (Slovakia), Natalia Skorodumova (Sweden), Dalimil Snita (Czech), N. Stepanov (Russia), D. Todorović (Serbia), M. Trtica (Serbia), V. Vasić (Serbia), D. Veselinović (Serbia), M. Zdujić (Serbia)

Local Executive Committee

Chairman: B. Adnadjević

Vice-chairman: S. Blagojević

Members: A. Abu Rabi-Stanković, P. Banković, N. Begović, J. Dostanić, A. Đerić, A. Ignjatović, A. Ivanović, A. Jović, D. Lončarević, J. Krstić, J. Maksimović, M. Milenković, S. Mačešić, V. Marković, B. Nedić, M. Petković, A. Popović-Bjelić, N. Potkonjak, D. Ranković, M. Stević, I. Stojković, M. Žunić

Honorary Committee

Nikola Hajdin,
Branko Kovačević,
Radivoje Mitrović
Vera Dondur

Vladimir Bumbaširević

Paula Putanov,
Momčilo Ristić,
Ankica Antić Jovanović,

President of Serbian Academy of Sciences and Arts
Rector of University of Belgrade
Secretary of State for Education and Science
President of the Council for Scientific and
Technological Development
Dean of the Faculty of Medicine of University of
Belgrade
Member of Serbian Academy of Sciences and Arts
Member of Serbian Academy of Sciences and Arts
Honorary President of Society of Physical Chemists of
Serbia

ROTATIONAL ISOMERS, NMR AND REACTIVITY ANALYSES OF TERT-PENTYL-1-BENZYL-4-METHYL-2,6-GLUTARIMIDE-3-CARBOXYLATE (PBMG)

V D. Vitnik,¹ J B. Popović-Đorđević,² Ž J. Vitnik,³
M D. Ivanović³, I. O. Juranić¹

¹ Department of Chemistry, IChTM, University of Belgrade, Belgrade, Serbia

² Faculty of Agriculture, University of Belgrade, Belgrade, Serbia

³ Faculty of Chemistry, University of Belgrade, Belgrade, Serbia

Abstract

The stable isomers of *tert*-pentyl-1-benzyl-4-methyl-glutarimide-3-carboxylate (PBMG) are determined. Optimized geometrical parameters of the stable isomers of PBMG are predicted by DFT/B3LYP/6-311++G(d,p). The ¹H and ¹³C nuclear magnetic resonance (NMR) chemical shifts of the molecules were calculated using the GIAO method and conform to the experimental values. To assess chemical reactivity of the molecule, the molecular electrostatic potential (MEP) surface map is plotted over the optimized geometry of the molecule.

Introduction

Glutarimide derivatives are recognized as anticonvulsants [1], sedatives [2] and antitumor agents [3]. Substituted imide rings are also found in some of the pharmacologically active natural products, such as cycloheximide [4], sesbanimide [5], izomigrastatine and migrastatine [6]. Numerous compounds containing six-membered imide ring are synthesized and express different pharmacological activities such as anxiolytic (buspirone, tandospirone) [2], immunosuppressive (thalidomide and analogues) [2], and others.

In our previously reported research PBMG has been synthesized by a novel tandem process [7].

The understanding of molecular geometry and reactive electrophilic and nucleophilic sites of the compound are of particular importance in the synthesis of a drug. Hence, in the present study, the energies and stability of different possible rotational isomers of PBMG are determined, as well as ¹H and ¹³C nuclear magnetic resonance (NMR) spectra for the title molecule are analyzed. Moreover, molecular electrostatic potential (MEP) surface is plotted over the optimized geometry to elucidate the reactivity of PBMG molecule.

Experimental

The chemical structure and purity of the synthesized compound was confirmed by its melting point, ¹H and ¹³C NMR and FT-IR spectra. ¹H and ¹³C NMR spectra were recorded in CDCl₃ using TMS as an internal standard on a Varian Gemini 200 spectrometer at 200 MHz and 50 MHz, respectively.

Computational details

All the calculations were performed using Gaussian 03 program package. Geometry of PBMG was fully optimized with DFT/B3LYP/6-311++G(d,p) method. ^1H and ^{13}C NMR chemical shifts are calculated with GIAO approach by applying B3LYP/6-311++G(d,p) method. To investigate the reactive sites of the title compound, the molecular electrostatic potentials for the 0.002 a.u. isosurfaces of electron density was evaluated using the B3LYP/6-311++G(d,p) method.

Results and Discussions

Molecular geometry

The molecule of PBMG could exist as *trans* and *cis* diastereomers. To find most stable isomer geometry, the DFT energy calculation is performed on the eight possible isomers of PBMG. The possible isomer geometry is found by locating -CH₂Ph group of the title molecule in two different orientations, above and below the plane of the glutarimide ring. The positioning of *tert*-pentyl group in two different orientations gives rise to eight isomers. The DFT energy calculation and the statistical Boltzmann distribution (shown in Table 1) reveal that, the isomers **1d** and **1h** acquire dominant stability among others *trans* (**1a-1d**) and *cis* (**1e-1h**) isomers, respectively. The optimized molecular structure of the most stable isomer **1d** of the title molecule is shown in Fig. 1.




Figure 1. Geometry optimized structure of **1d** isomer of PBMG.

Table 1. The energies and the statistical Boltzmann distribution weighted values of the most stable PBMG isomers.

Isomer	Energy / a.u.	%
1a	-1095.0792247	11.22
1b	-1095.0801943	31.15
1c	-1095.0793179	12.38
1d	-1095.0804938	42.69
1e	-1095.0750486	0.14
1f	-1095.0769821	1.06
1g	-1095.0750302	0.14
1h	-1095.0771161	1.22

NMR spectra

From experimental data the ratio between *trans/cis* diastereomers (5:1) is determined by ratio of the two signals at 3.08 and 3.45 ppm (two doublets from atom H(24) on C(3)). The same signal ratios are found at 1.26 (1.34) ppm from two singlets of methyl groups at C(11) and C(12) in *trans/cis* isomers, as well as for signals at 0.94 (0.98) ppm from methyl groups on C(7) and at 0.68 (0.77) ppm from methyl groups on C(14). Doublets at 2.62 (2.70) ppm mark methylene of

B-06-P

glutarimide ring, while quartets at 2.20 (2.45) and 1.66 (1.68) ppm mark atom H(25) on C(4) and methylene hydrogens on C(13), respectively. Singlet at 4.82 (4.84) ppm mark methylene hydrogens on C(15). Peaks at 7.05-7.26 ppm indicate the presence of aromatic hydrogen atoms. In ^{13}C NMR spectra signals for carbonyl groups are at 170.54 (171.29), 168.54 (168.79) and 167.03 (165.83) ppm in *trans/cis* isomers. The peaks appearing in the range 138.62–127.09 ppm are assigned to the carbon atoms of phenyl group. The signals at 84.76 (85.43) is assigned to C(10) carbon. The aliphatic CH₂ (C(6), C(13), and C(15)) carbons are observed at 42.55, 38.24 (36.62) and 57.55 (55.04) ppm, respectively. The carbon atoms of the CH groups belonging to the glutarimide ring are observed at 33.09 and 26.27 (27.4) ppm, for C(3), and C(4), respectively. Methyl groups are at 24.87 (24.95), 18.69 (17.03) and 7.74 ppm, for C(11, 12), C(7), and C(14), respectively. The theoretical ^1H and ^{13}C chemical shift results for PBMG are generally close to the experimental ^1H and ^{13}C shift data.

MEP

The MEP map shows that the negative potential sites are on electronegative O atoms of the carbonyl groups and the positive potential sites are around the carbon C(4) of the glutarimide ring. These sites give information about the region from where the compound can make intermolecular interactions. Thus, it would be predicted that the carbonyl groups fragment of PBMG will be the most reactive site for electrophilic, and C(4) atom for nucleophilic attack.

Acknowledgments

This work has been financially supported by Ministry of Education and Science, Republic of Serbia, under Grant No. 172035, 172032 and by High-Performance Computing Infrastructure for South East Europe's Research Communities European project.

References

- [1] S. A. Dandekar, S. N. Greenwood, T. D. Greenwood, S. Mabic, J. S. Merola, J. M. Tanko, , J. F. Wolfe, *J. Org. Chem.*, 1999, 64, 1543-1553.
- [2] The Merck Index, 13th Ed; Merck & Co., Inc., NJ, 2001.
- [3] D. K. K. Ha, W. H. Lau, *Cancer Lett.*, 1988, 41, 217-224.
- [4] C. Christner, R. Wyrwa, S. Marsch, G. Küllertz, R. Thiericke, S. Grabley, D. Schumann, G. Fischer, *J. Med. Chem.*, 1999, 42, 3615-3622.
- [5] F. Matsuda, S. Terashima, *Tetrahedron*, 1988, 44, 4721-4736.
- [6] J. Ju, S. R. Rajski, S.-K. Lim, J.-W. Seo, N. R. Peters, F. M. Hoffmann, B. Shen, *Bioorg. Med. Chem. Lett.*, 2008, 18, 5951-5954.
- [7] J. B. Popović-Đorđević, M. D. Ivanović, V. D. Kiricojević, *Tetrahedron Lett.*, 2005, 46, 2611–2614.

CIP Volume I

CIP - Каталогизација у публикацији
Народна библиотека Србије, Београд

544(082)
621.35(082)
66.017/.018(082)

MEĐUNARODNA konferencija iz fundamentalne i
primenjene fizičke hemije (11 ; 2012 ;
Beograd)

Physical Chemistry 2012 : proceedings.
#Vol. #1 / 11th International Conference on
Fundamental and Applied Aspects of Physical
Chemistry, September 24-28, 2012, Belgrade ;
[editors S.[Slobodan] Anić and Ž.[Željko]
Čupić ; organized by Society of Physical
Chemists of Serbia ... et al.]. - Belgrade :
Society of Physical Chemists of Serbia, 2012
(Belgrade : Jovan). - VI, 498 str. : ilustr.
; 24 cm

"The Conference is dedicated to Professor
Ivan Draganić" --> nasl. str. - Tiraž 200. -
Bibliografija uz svaki rad.

ISBN 978-86-82475-27-9

1. Društvo fizikohemičara Srbije (Beograd)
a) Физичка хемија - Зборници b)
Електрохемијско инжењерство - Зборници c)
Наука о материјалима - Зборници
COBISS.SR-ID 193432332