

Supported by



MICROBIOLOGIA BALKANICA 2011

7th BALKAN CONGRESS OF MICROBIOLOGY

8th CONGRESS OF SERBIAN MICROBIOLOGISTS

Serbian Society for Medical Microbiology

Serbian Society for Microbiology

Faculty of Biology, University of Belgrade



October 25-29, 2011

Continental Hotel, Belgrade, SERBIA

CIP – Katalogizacija u publikaciji
Narodna biblioteka Srbije, Beograd

Udruženje mikrobiologa Srbije, Beograd.

Knjiga radova (Elektronski izvor) – Proceedings / MICROBIOLOGIA BALKANICA 2011 - 7th BALKAN CONGRESS OF MICROBIOLOGY & 8th CONGRESS OF SERBIAN MICROBIOLOGISTS, 25-29. oktobar 2011; (organizator) Udruženje mikrobiologa Srbije, Udruženje medicinskih mikrobiologa Srbije; (urednici: Dragojlo Obradović, Lazar Ranin, Špiro Radulović) – Beograd
1 elektronski optički disk (CD-ROM); 12cm

Sistemske zahteve: Nisu navedeni. Nasl. sa naslovnog ekrana. –

Radovi na engleskom jeziku. –

Tekst latinica. –

Tiraž – 600.

Abstracts. – Registar

ISBN 978-86-914897-0-01

Udruženje mikrobiologa Srbije, Beograd.

KNJIGA RADOVA / PROCEEDINGS

MICROBIOLOGIA BALKANICA 2011 - 7th BALKAN CONGRESS OF MICROBIOLOGY & 8th CONGRESS OF SERBIAN MICROBIOLOGISTS, 25-29. oktobar 2011

Izdaje / Published by:

Udruženje mikrobiologa Srbije

Nemanjina 6, 11 080 Beograd, Srbija, tel/fax: 011 2199 711, dobradovic@agrifaculty.bg.ac.rs

Za izdavača / For Publisher:

Dragojlo Obradović, predsednik Udruženja

Urednici/Editors:

Dragojlo Obradović

Lazar Ranin

Špiro Radulović

ISBN 978-86-914897-0-01

Kompjuterska obrada teksta / Computer Layout:

Jelena Zovko Belić

Tiraž / Circulation

600 primeraka / 600 copy

Umnožavanje / Copying

Megaphone d.o.o., Vladimira Rolovića 105, Beograd, Srbija

Committees

Organizing Committee

President: Prof. dr Spiro Radulovic

Vice president: Prof. dr Dragojlo Obradovic

General Secretary: Prof. dr Lazar Ranin

Members:

Prof. dr Branislava Kocic
 Prof. dr Branislava Savic
 Prof. dr Ruzica Asanin
 Prof. dr Veljko Mirovic
 Prof. dr Vaso Taleski
 Prof. dr Djordje Fira
 Prof. dr Slavisa Stankovic

Scientific Committee

President: Prof. dr Tanja Jovanovic

Members:

Prof. dr Ruzica Asanin
 Prof. dr Dejan Baskic
 Prof. dr Sava Buncic
 Prof. dr Angel Galabov
 Prof. dr Zeynep Gulay
 Dr Dobrila Jakic Dimic
 Prof. dr Dragutin Djukic
 Prof. dr Mirjana Jarak
 Prof. dr Vera Katic
 Prof. dr Jelena Knezevic Vukcevic
 Prof. dr Dejan Krnjajic
 Prof. dr Marija Kulauzov
 Prof. dr Zoran Kulisic
 Prof. dr Nada Kuljić Kapulica
 Prof. dr Dusan Lalosevic
 Prof. dr Nenad Milic
 Prof. dr Gordana Mijovic
 Prof. dr Sanja Mitrovic
 Dr Jasmina Nedeljkovic
 Prof. dr Viktor Nedovic
 Prof. dr Marian Negut
 Prof. dr Miomir Niksic
 Prof. dr Aleksa Obradovic
 Prof. dr Anna Papa
 Prof. dr Milena Petrovska
 Prof. dr Olga Petrović
 Dr Milanko Sekler
 Prof. dr Marija Skrinjar
 Prof. dr Vaso Taleski
 Prof. dr Ljubisa Topisirovic
 Dr Branka Vasiljevic
 Dr Branka Vidic
 Prof. dr Jelena Vukojevic
 Prof. dr Miroslav Vrvic
 Dr Milan Kojić
 Prof. dr Siniša Markov
 Prof. dr Vera Raičević

Board of the Balkan Society for Microbiology

President:

Prof. dr Milena Petrovska (Skopje)

President – elect:

Prof. dr Spiro Radulovic (Belgrade)

Members:

Bulgaria:

Prof. dr Angel GALABOV (Sofia)
 Prof. dr Todor KANTARDJIEV (Sofia)
 Prof. dr Hristo NAJDENSKI (Sofia)

Greece:

Prof. dr Athanassios TSAKRIS (Athens)
 Prof. dr Anna PAPA KONIDARI (Thessaloniki)

Macedonia:

Prof. dr Dzoko KUNGULOVSki (Skopje)
 Prof. dr Nikola PANOVSki (Skopje)

Montenegro:

Dr Jelena ZINDOVIC (Podgorica)
 Dr Zoran VRATNICA (Podgorica)
 Doc. dr Vineta VUKSANOVIC (Podgorica)

Romania:

Prof. dr Gabriel IONESCU (Bucharest)
 Prof. dr Marian NEGUT (Bucharest)
 Prof. dr Alexandru RAFILA (Bucharest)

Serbia:

Prof. dr Branislava KOCIC (Nis)
 Prof. dr Lazar RANIN (Belgrade)

Turkey:

Prof. dr Nezahat GURLER (Istanbul)
 Prof. dr Zeynep GULAY (Istanbul)

Plenary speakers

Milton da Costa (Portugal)
 Otto Haller (Germany)
 Mirsada Hukic (BIH)
 Gunnar Kahlmeter (Germany)
 Anna Papa – Konidari (Greece)
 Elisabeth Nagy (Hungary)
 Bernhard Schink (Germany)
 Vittorio Venturi (Italy)

MALDI-TOF analysis of rhamnolipids produced by *P. aeruginosa* NCAIM (P) B001380

Rikalovic G. Milena, Gordana Gojic-Cvijovic^I, Miroslav M. Vrvic,
Ivanka Karadzic^{II},

Faculty of Chemistry, University of Belgrade

^IICTM, University of Belgrade

^{II}Faculty of Medicine, University of Belgrade

Department of Biochemistry, ^IDepartment of Chemistry,

^{II}Department of Chemistry

Belgrade, Serbia

Introduction

Rhamnolipids (RLs) are low-molecular-mass biosurfactants. The principal RLs are mono-rhamno-di-lipidic and di-rhamno-di-lipidic congeners, while mono-rhamno-mono-lipidic and di-rhamno-mono-lipidic congeners are biosynthesized only under certain cultivation conditions (Thazbi *et al.*, 2004).

Aim

MALDI-TOF analysis of RLs produced by strain *P. aeruginosa* NCAIM (P) B001380 on Lurie-Bertani (LB) medium.

Material and methods

Strain *P. aeruginosa* NCAIM (P) B001380 was isolated from industrial mineral metal-cutting oil (Karadzic *et al.*, 2004). The strain was activated on nutrient agar at 30 °C for 24 h and fermentation was carried in LB medium, at 30 °C for seven days, at 250 cycles/min (Karadzic *et al.*, 2004).

Crude preparation of RL was obtained by acidic precipitation, followed by extraction of chloroform: methanol (2:1) (Heyd *et al.*, 2008). Concentration of RL was determined spectrophotometrically at 421 nm, with orcinol reaction using rhamnose as a standard (Wilhelm *et al.*, 2007).

Isolated RL mixture was analyzed by MALDI-TOF. Mass spectra were acquired on a Voyager Biospectrometry DE Pro Workstation (Perseptive Biosystems, Framingham, MA, USA), using matrix solution of 2, 5-dihydroxybenzoic acid, concentration of 0.5 M in methanol containing 0.1% trifluoroacetic acid (TFA) (Schiller *et al.*, 2004).

Results

Composition of RL mixture on LB medium was monitored for seven days. Total number of RL congeners was: 6 (5 mono-RLs, 1 di-RLs), 7 (5 mono-RLs, 2 di-RLs), 6 (5 mono-RLs, 1 di-RLs), 4 (3 mono-RLs, 1 di-RLs), 2 (1 mono-RLs, 1 di-RLs), 2 (2 mono-RLs, 0 di-RLs), 4 (3 mono-RLs, 1 di-RLs), from the first to the seventh day, respectively. Total number of mono-RL congeners rapidly declined after the third day of fermentation. The first three days of fermentation mono-RLs that dominated were Rha-C8-C10:1, Rha-C10-C12:1, Rha-C10-C14:1 and Rha-C10-C14, while after that period Rha-C8-C10:1 and Rha-C16-C16 were detected. Rha-C8-C8, Rha-C10-C10 and Rha-C10-C10-CH₃ were sporadically occurring. Number of di-RLs practically didn't change, but different structure of RL congeners were present, depending on day.

Conclusion

RL profile significantly varied during fermentation on LB medium. The first three days of fermentation mono-RLs contribute to diversity of RLs mixture, after that number of different congeners decline till the end of fermentation.

Key words: *P. aeruginosa*, rhamnolipids, MALDI-TOF

MALDI-TOF analysis of rhamnolipids produced by *Pseudomonas aeruginosa* NCAIM (P) B001380

Rikalović G. Milena, Gordana Đ. Gojčić-Cvijović¹, Miroslav M. Vrvic, Ivanka M. Karadžić^{1,2}

Faculty of Chemistry, University of Belgrade, Studentski trg 12-16, Belgrade, Serbia, ¹ICTM Department of Chemistry, University of Belgrade, Njegoseva 12, Belgrade, Serbia, ²Faculty of Medicine, Department of Chemistry, University of Belgrade, Visegradska 26, Belgrade, Serbia

This study represents MALDI-TOF analysis of rhamnolipids (RLs) produced by strain *P. aeruginosa* NCAIM (P) B001380. Strain was grown on Lurie-Bertani (LB) medium. Composition of rhamnolipid (RL) mixture on Lurie-Bertani medium was monitored for seven days. Total number of RL congeners produced from the first to the seventh day was: 6 (5 mono-RLs, 1 di-RLs), 7 (5 mono-RLs, 2 di-RLs), 6 (5 mono-RLs, 1 di-RLs), 4 (3 mono-RLs, 1 di-RLs), 2 (1 mono-RLs, 1 di-RLs), 2 (2 mono-RLs, 0 di-RLs), 4 (3 mono-RLs, 1 di-RLs). Total number of mono-RL congeners rapidly declined after the third day of fermentation. Number of di-RLs practically didn't change, but different structure of RL congeners were present, depending on day of growth. RL profile significantly varied during fermentation on LB medium. The first three days of fermentation mono-RLs contribute to diversity of RLs mixture, after that number of different congeners decline till the end of fermentation.

Introduction

Rhamnolipids are low-molecular-mass biosurfactants. The principal RLs are mono-rhamno-di-lipidic and di-rhamno-di-lipidic congeners, while mono-rhamno-mono-lipidic and di-rhamno-mono-lipidic congeners are biosynthesized only under certain cultivation conditions (Thazbi *et al.*, 2004). RLs are microbial secondary metabolites (Van Hamme *et al.*, 2006) and their production is possible from most carbon sources supporting bacterial growth. (Tahzibi *et al.*, 2004, Soberón-Chávez *et al.*, 2005, Chayabutra *et al.*, 2001).

During the last a few decades, increasing interest in biological surfactants led to an intensification of research for their cost-efficient production. Also, search for alternative rhamnolipid-producing strains makes a structure analysis and constant adaptation of the existing analytical methods necessary. A great success of Mass Spectrometry (MS) in biological sciences and an intensive development of lipidomics give rise to development of MS analytical platform for RL high throughput investigation.

Dynamic of RL production by *P. aeruginosa* NCAIM (P) B001380 was analyzed using a recently reported MALDI-TOF based method. MALDI-TOF-MS has been developed as a novel and convenient technique for rapid high-throughput screening of RL-containing samples and RL-producing *Pseudomonas* spp. MALDI-TOF-MS assignments have been successfully verified by GC/MS, HPLC, and NMR spectroscopy, confirming the validity of the MALDI-TOF mass spectrometry method (Hayd *et al.*, 2008, Price *et al.*, 2009).

Material and methods

Strain *P. aeruginosa* NCAIM (P) B001380 was isolated from industrial mineral metal-cutting oil (Karadzic *et al.*, 2004).

The strain was activated on nutrient agar at 30 °C for 24 h and fermentation was carried in LB medium, at 30 °C for seven days, at 250 cycles/min (Karadzic *et al.*, 2004).

Crude preparation of RL was obtained by acidic precipitation followed by extraction of chloroform: methanol (2:1) (Heyd *et al.*, 2008).

Concentration of RL (Crl) was determined spectrophotometrically at 421 nm, with orcinol reaction using rhamnose as a standard (Wilhelm *et al.*, 2007).

Isolated RL mixture was analyzed by MALDI-TOF. Mass spectra were acquired on a Voyager Biospectrometry DE Pro Workstation (PerSeptive Biosystems, Framingham, MA, USA), using matrix solution of 2,5-dihydroxybenzoic acid, concentration of 0.5 M in methanol containing 0.1% trifluoroacetic acid (TFA) (Schiller *et al.*, 2004).

Results

Composition of RL mixture on LB medium was monitored for seven days. Total number of RL congeners was: 6 (5 mono-RLs, 1 di-RLs), 7 (5 mono-RLs, 2 di-RLs), 6 (5 mono-RLs, 1 di-RLs), 4 (3 mono-RLs, 1 di-RLs), 2 (1 mono-RLs, 1 di-RLs), 2 (2 mono-RLs, 0 di-RLs), 4 (3 mono-RLs, 1 di-RLs), from the first to the seventh day, respectively. Total number of mono-RL congeners rapidly declined after the third day of fermentation. The first three days of fermentation mono-RLs that dominated were Rha-C8-C10:1, Rha-C10-C12:1, Rha-C10-C14:1 and Rha-C10-C14, while after that period Rha-C8-C10:1 and Rha-C16-C16 were detected. Rha-C8-C8, Rha-C10-C10 and Rha-C10-C10-CH3 were sporadically occurring. Number of di-RLs practically didn't change, but different structure of RL congeners were present, depending on day of growth.




Figure 1. Dynamic of RL production on LB medium by *P. aeruginosa* NCAIM (P) B001380: a) the first, b) the second, c) the third, d) the fourth, e) the fifth, f) the sixth, g) the seventh day of fermentation.

Table 1. Dynamic of rhamnolipid production of *P. aeruginosa* NCAIM (P) B001380 on LB medium: profile of RLs and concentrations (Crl)

Day	Mono-RL congener Rha-	Di-RL congener Rha-Rha-	Crl mg/l
1	C8-C10:1/C10:1-C8, C10-C12:1/C12:1-C10, C10-C14/C14-C10//C12-C12, C10-C14:1/C14:1-C10//C12-C12:1/C12:1-C12, C10-C10-CH3	C14-C14	51
2	C10-C10, C10-C12/C12-C10, C10-C12:1/C12:1, C10-C14/C14-C10//C12-C12, C10-C14:1/C14:1-C10//C12-C12:1/C12:1-C12, C10-C10-CH3	C12-C12, C12-C14/C14-C12	53
3	C8-C8, C8-C10:1/C10:1-C8, C10-C14/C14-C10//C12-C12, C10-C14:1/C14:1-C10//C12-C12:1/C12:1-C12, C16-C16	C12-C12	74
4	C8-C10:1/C10:1-C8, C14-C16/C16-C14, C16-C16	C10-C12:1/C12:1-C10	57
5	C8-C10:1/C10:1-C8	C10-C12:1/C12:1-C10	80
6	C10-C10, C10-C10-CH3	/	96
7	C8-C8, C8-C10:1/C10:1-C8, C16-C16	C10-C12:1/C12:1-C10	53

During fermentation Crl of RLs was varying in a range from 50 to 100 mg/l. Two maxima of rhamnolipid production were obtained (Table 1): the first maximum at the third day (74 mg/l) and the second maximum between the fifth and the sixth day (80-96 mg/l). Days with the highest production of RLs showed different diversity of RL mixture (3th, 5th and 6th day had 6, 2, 2 different RL congeners, respectively).

Conclusion

RL profile significantly varied during growth of *P. aeruginosa* NCAIM (P) B001380 on LB medium.

For the first three days of bacterial growth mono-RLs significantly contributed to diversity of RLs mixture, after that their number declined till the end of fermentation. Number of di-RLs was low and practically didn't change, albeit different structure of RL congeners were present, depending on the phase of growth. The highest diversity of RL mixture was at the second day of fermentation (7 different congeners). The highest Crl were obtained after 72 h (74 mg/l) and between fifth and sixth day of fermentation (80-96 mg/l).

Acknowledgment

This work was supported by the project III 43004 of the Ministry of Science and Technological Development of Serbia.

The authors are grateful to Dr. N. Fujiwara from the Institute for Technological Research (TRI), Osaka, Japan for his kind donation of strain and to Dr. M. Petković from the Institute of Nuclear Sciences Vinca, Belgrade, Serbia for MALDI-TOF analysis.